Technical Note No. 23¹

Options, Futures, and Other Derivatives

John Hull

The Black, Derman, and Toy Model

As explained in the text, there are two types of model of the short rate: equilibrium and no-arbitrage models. In an equilibrium model the process followed by the short-term interest rate is specified. This totally defines the model. Zero-coupon bond prices and the term structure of interest rates are outputs from the model. Examples of equilibrium models are the Vasicek and Cox, Ingersoll, and Ross models. These models each have three parameters. The parameters can be chosen so that the models provide an approximate fit to the term structure of interest rates, but the fit is not usually an exact one.

A no-arbitrage model is constructed so that it is exactly consistent with the term structure of interest rates that is observed in the market. This means that the term structure of interest rates is an input to the model, not an output from it. No-arbitrage models can be constructed in many different ways. An early no-arbitrage model was the Black, Derman, and Toy model published in 1990.² This model has the advantage that it can easily be represented in the form of a binomial tree. To correspond as closely as possible with the Black-Derman-Toy paper, we assume that interest rates are compounded annually.

The Black-Derman-Toy model is a particular case of the more general Black-Karasinski model. The short rate follows a mean-reverting lognormal process. However, the way the tree is constructed implies a relationship between the short rate volatility and the reversion rate.

As in the case of the binomial tree used to value stock options, we consider steps of length Δt . The tree models the behavior of the Δt -period interest rate. The zero-coupon interest rates for all maturities at time zero are known. We denote the zero-coupon interest rate for a maturity of $n\Delta t$ by R_n . The volatility of the Δt rate between time $(n-1)\Delta t$ and $n\Delta t$ is denoted by σ_n .

During each time step the Δt -period interest rate has a 50% probability of moving up and a 50% probability of moving down. The tree is shown in Figure 1. The initial Δt period rate, *r*, is known and equals R_1 . The value of a zero-coupon bond that pays off \$1 at time $2\Delta t$ is

$$\frac{1}{\left(1+R_2\right)^{2\Delta t}}$$

The value of this bond at node B on the tree is

¹ ©Copyright John Hull. All Rights Reserved. This note may be reproduced for use in conjunction with Options, Futures, and Other Derivatives by John C. Hull

² See F. Black, E. Derman, and W. Toy, ``A one-factor model of interest rates and its application to Treasury bond options, Financial Analysts Journal, 46 (1), 33-39.

$$\frac{1}{\left(1+r_d\right)^{\Delta t}}$$

The value of the bond at node C is

$$\frac{1}{\left(1+r_{u}\right)^{\Delta t}}$$

It follows that the value of the bond at the initial node A is

$$\frac{1}{(1+r)^{\Delta t}} \left[0.5 \frac{1}{(1+r_d)^{\Delta t}} + 0.5 \frac{1}{(1+r_u)^{\Delta t}} \right]$$

Hence

$$\frac{1}{(1+r)^{\Delta t}} \left[0.5 \frac{1}{(1+r_d)^{\Delta t}} + 0.5 \frac{1}{(1+r_u)^{\Delta t}} \right] = \frac{1}{(1+R_2)^{2\Delta t}}$$
(1)

Figure 1 The Binomial Tree



This is one equation that must be satisfied by r_u and r_d . To match the volatility, the standard deviation of the logarithm of the interest rate at time Δt must be $\sigma_1 \sqrt{\Delta t}$ (Recall: σ_i is the volatility of interest rates during the *i*th time period.) This means that³

$$0.5\ln\frac{r_u}{r_d} = \sigma_1 \sqrt{\Delta t} \tag{2}$$

Equations (1) and (2) can be solved to determine r_u and r_d .

We now move on to determine r_{uu} , r_{ud} and r_{dd} . To match volatility we must have

$$0.5 \ln \frac{r_{uu}}{r_{ud}} = \sigma_2 \sqrt{\Delta t} \tag{3}$$

$$0.5 \ln \frac{r_{ud}}{r_{dd}} = \sigma_2 \sqrt{\Delta t} \tag{4}$$

We must also match the price of a zero-coupon bond that pays off \$1 at the end of time $3\Delta t$. Rolling back through the tree the values of this bond at nodes D, E, and F are

$$\frac{1}{(1+r_{ud})^{\Delta t}}$$
, $\frac{1}{(1+r_{ud})^{\Delta t}}$, and $\frac{1}{(1+r_{uu})^{\Delta t}}$

respectively. The values at nodes B and C are

$$\frac{1}{(1+r_d)^{\Delta t}} \left[0.5 \frac{1}{(1+r_{dd})^{\Delta t}} + 0.5 \frac{1}{(1+r_{ud})^{\Delta t}} \right]$$

and

$$\frac{1}{(1+r_u)^{\Delta t}} \left[0.5 \frac{1}{(1+r_{ud})^{\Delta t}} + 0.5 \frac{1}{(1+r_{uu})^{\Delta t}} \right]$$

The value at the initial node is therefore

³ To see this note that the variance of the logarithm of the interest rate is $0.5(\ln r_u)^2 + 0.5(\ln r_d)^2 - [0.5(\ln r_u + \ln r_d)]^2 = [0.5(\ln r_u - \ln r_d)]^2$

$$\frac{1}{(1+r)^{\Delta t}} \left(0.5 \frac{1}{(1+r_d)^{\Delta t}} \left[0.5 \frac{1}{(1+r_{dd})^{\Delta t}} + 0.5 \frac{1}{(1+r_{ud})^{\Delta t}} \right] \right) + \frac{1}{(1+r)^{\Delta t}} \left(0.5 \frac{1}{(1+r_u)^{\Delta t}} \left[0.5 \frac{1}{(1+r_{ud})^{\Delta t}} + 0.5 \frac{1}{(1+r_{uu})^{\Delta t}} \right] \right) = \frac{1}{(1+R_3)^{3\Delta t}}$$
(5)

The interest rates r, r_u and r_d have already been determined. Equations (3), (4), and (5) therefore provide three equations for determining r_{uu} , r_{ud} , and r_{dd} .

Continuing in this way a complete tree can be constructed. The calculations are made considerably easier if as we move forward we keep track of the value of a security that pays \$1 if a particular node is reached and zero elsewhere. It is then only necessary to roll back one step when valuing zero-coupon bonds using the tree.

Determining the σ_i

The determination of the σ_i depends on the data available. Sometimes the historical volatilities of zerocoupon bond yields are used; sometimes the volatilities of caps or swaptions are used. An iterative search procedure is always necessary.

When bond yields are being matched, we assume that we have data at time zero on the volatilities of a bond maturing at time $i\Delta t$. We will denote this by β_i . (We approximate β_i as the volatility of this bond yield between time zero and time Δt .) We denote y_{un} as the yield on a bond maturing at time $n\Delta t$ at node C and y_{dn} as the yield on a bond maturing at time $n\Delta t$ at node B. Considering a bond that matures at time $2\Delta t$,

$$\beta_2 \sqrt{\Delta t} = 0.5 \ln \frac{y_{u2}}{y_{d2}}$$

Because there is only one period left in the bond's life at the nodes at time Δt , $y_{u2} = r_u$ and $y_{d2} = r_d$. As a result

$$\beta_2 \sqrt{\Delta t} = 0.5 \ln \frac{r_u}{r_d}$$

Using equation (2) leads to

$$\sigma_1 = \beta_2$$

The interest rates r_u and r_d can then be determined from equations (1) and (2).

Determining the subsequent σ_i requires an iterative search. For example, to determine σ_2

$$\beta_3 \sqrt{\Delta t} = 0.5 \ln \frac{y_{u3}}{y_{d3}}$$

This must be solved iteratively with equations (3), (4), and (5) for σ_2 , r_{uu} , r_{ud} , and r_{dd} .

In general the procedure to determine σ_i (*i* > 1) is

- 1. Choose a trial value of σ_i
- 2. Calculate the interest rates at time $i\Delta t$
- 3. Calculate the yield volatility for a bond lasting until $i\Delta t$ from the tree. This involves calculating the bond yields y_u and y_d at nodes B and C. The bond yield volatility is $0.5\ln(y_u/y_d)$
- 4. Search iteratively for the value of σ_i that matches the bond yield volatility

Once the tree has been constructed it can be used to value a range of interest rate derivatives.

Example

As an example of the application of the model suppose that the term structure of interest rates is as shown in Table 1, the zero-coupon yield volatilities are as shown in Table 2, and the time step is one year. In this case r=0.10, $\Delta t=1$, $\sigma_1=0.19$ (the two-year yield volatility) and equations (1) and (2) give

$$\frac{1}{1.10} \left[0.5 \frac{1}{1+r_d} + 0.5 \frac{1}{1+r_u} \right] = \frac{1}{1.11^2}$$
$$0.5 \ln \frac{r_u}{r_d} = 0.19$$

Solving these two equations gives $r_u = 0.1432$ and $r_d = 0.0979$.

Table 1

Zero-coupon Yield Curve (Annually Compounded)

Maturity (years)	Rate
1	10.0
2	11.0
3	12.0
4	12.5
5	13.0

Table 2

Yield Volatilities

i	β_i
2	19.0%
3	18.0%
4	17.5%
5	16.0%

Equations (3), (4), and (5) give

$$0.5\ln\frac{r_{uu}}{r_{ud}} = \sigma_2$$

$$0.5\ln\frac{r_{ud}}{r_{dd}} = \sigma_2$$

$$\frac{1}{1.1} \left(0.5 \frac{1}{1.0979} \left[0.5 \frac{1}{1+r_{dd}} + 0.5 \frac{1}{1+r_{ud}} \right] \right) + \frac{1}{1.1} \left(0.5 \frac{1}{1.1432} \left[0.5 \frac{1}{1+r_{ud}} + 0.5 \frac{1}{1+r_{uu}} \right] \right) = \frac{1}{1.12^3}$$

We do not know σ_2 directly. For each trial value of σ_2 we solve equations (3), (4), and (5) and then calculate the price of a three-year bond at nodes B and C. The price of a three-year bond at node B is

$$B_d = \frac{1}{1.0979} \left[0.5 \frac{1}{1 + r_{dd}} + 0.5 \frac{1}{1 + r_{ud}} \right]$$

and the bond yield at node B is

$$y_d = \sqrt{\frac{1}{B_d}} - 1$$

The price of a three-year bond at node C is

$$B_u = \frac{1}{1.1432} \left[0.5 \frac{1}{1 + r_{ud}} + 0.5 \frac{1}{1 + r_{uu}} \right]$$

_

The bond yield is

$$y_u = \sqrt{\frac{1}{B_u}} - 1$$

Carrying out an iterative search we find that $\sigma_2 = 0.172$ does the trick. With this value of σ_2 the solutions to the three equations are

$$r_{uu} = 0.1942$$

 $r_{ud} = 0.1377$
 $r_{dd} = 0.0976$

These in turn give $B_u = 0.7507$, $B_d = 0.8152$, $y_u = 0.1542$, and $y_d = 0.1076$. Because $0.5\ln(0.1542/0.1076) = 0.18$ the three-year yield volatility is matched.

The complete tree of short rates is shown in Figure 2.



